

## Chapter 9: Crop Rotations Can Increase Corn Profitability and Reduce Pests



Peter Sexton (Peter.Sexton@sdstate.edu)

This chapter provides a brief overview on how rotations can increase long-term sustainability and resilience against climate variability for South Dakota producers. Crop rotation is a complex subject where biological factors, farm management resources, and market forces all interact to influence rotation effectiveness.

### Introduction

Crop rotations are long-term plans that improve sustainability and profitability. A producer considering crop rotation should examine:

- Profitability, equipment, and labor availability.
- Climate and market variability.
- Soil health.
- Short-term gains vs. long-term sustainability.
- The impact on weed, insect, and disease problems.
- Pest resistance to various control mechanisms.
- Matching crop production requirements with available resources.

Crop rotations are a foundational element of sustainability and long-term profitability. For example, the introduction of the “Norfolk Rotation” (Barley-Clover/ryegrass-Wheat-Turnips) by Sir Charles Townshend in England played a large role in nearly tripling England’s agriculture output in the 1700s in a sustainable manner. This technology improvement provided food and the labor required for England’s Industrial Revolution. Opposite results can occur if the production systems adopt extractive rather than sustainable techniques. For example, it is thought that the ancient inhabitants of Easter Island deforested their island leading to soil erosion, a loss of productivity, and societal collapse. Although in a different environment, similar loss of soil resources occurred in the Mediterranean 1500 years ago (Thirgood, 1981).

One way to consider sustainable production systems is to look at natural systems as a model to mimic. Natural systems tend to maximize resource capture and biomass production while minimizing nutrient loss. Natural systems keep the soil covered and protect the soil from erosion. As natural systems develop, they follow a “succession” process where one set of species modifies the environment to the benefit of the next set of species. In a similar manner, a good rotation program should be productive, minimize nutrient loss, cover the soil, provide resilience against pests and stress, and each crop should benefit of the next crop.

## Designing a Rotation

Rotations should be adaptable to local conditions and challenges. There are many factors that must be considered when designing a rotation. Producers need to look at rotations as one tool for optimizing long-term profitability and reducing risk. Achieving these goals is complicated, as one management practice may have negative implications on other practices. For example, reducing tillage intensity without use of a sustainable rotation can increase the risk of plant diseases (Table 9.1).

**Table 9.1 Some corn diseases that can be influenced by rotations.**

Disease	Pathogen and Environment	Inoculum source
Goss's wilt	<i>Clavibacter michiganensis</i> (bacteria); associated with injury from violent weather (e.g., wind and hail); favored by moderate temperatures, and can overwinter on some weeds.	Overwinters on residue; also some grassy weeds act as alternate hosts; moves with rain.
Gray Leaf Spot	<i>Cercospora zeaе-maydis</i> (fungi); favored by moderate to warm temperatures and high humidity.	Overwinters on residue; moves with wind and rain.
Anthrachnose leaf blight and stalk rot	<i>Colletotrichum graminicola</i> (fungi); favored by warm temperatures and long periods of cloudy, humid weather.	Overwinters on residue; moves with wind and rain.
Eyespot	<i>Kabatiella zeaе</i> (fungi); favored by cool, wet weather.	Overwinters on residue; moves with wind and rain.
Northern Corn Leaf Blight	<i>Exserohilum turcicum</i> (fungi); favored by moderate temperatures and humid weather.	Overwinters on residue; moves with wind.

### Rotations and Plant Diseases

Rotation is a very valuable tool for breaking disease cycles, particularly in no-till and conservation tillage systems. Crop residue acts as an inoculum source for many important diseases in corn (Table 9.1). Hence, rotations that use nonhost crops or resistant hybrids/varieties provide an opportunity for the residues to decompose, which should decrease pest risks.

Because certain pests persist in the soil, there are some diseases, such as seedling damping off (*Pythium* spp) and root rots (*Rhizoctonia solani* and *Fusarium* spp), that can be managed only by combining the rotation with other techniques. Additional methods might include using appropriate seed treatments, delaying seeding, and installing tile drainage.

### Rotations and Weed Management

Rotation can have large impacts on weed pressure. Rotations provide the opportunity to rotate the herbicide mode of action, which should reduce the risk of creating herbicide-resistant weeds. A “stacked” rotation can be effective in reducing this risk. In a stacked rotation, the same or very similar crops are grown two years in a row and then skipped for four or more years (e.g., corn-corn-soybean-soybean-wheat-wheat), allowing for the use of herbicides with long residuals in the first year of each crop while maintaining a long period (four years) where the land is rotated to other crops (Beck, 2003). Alfalfa can also be used for this purpose.

Similarly, an advantage can be gained by a rotation between warm- and cool-season crops, where each cycle is held for two seasons (two warm-season crops followed by two cool-season crops) (Anderson, 2008). Holding the given pattern for two years disrupts weed life cycles such that the weed seeds have to survive for three years




Figure 9.1 Average weed density after 10 years of different sequences of warm-season (w) and cool-season (c) crops from three trials. The lowest weed density was found where the warm- and cool-season crops were each grown in two-year blocks – two years of cool-season crops followed by two years of warm-season crops (e.g., oats-wheat-corn-soybeans; or wheat-canola-sorghum-sunflower). (Modified from Anderson, 2008)

before they get the opportunity to grow and multiply (Fig. 9.1).

### **Rotation, Residue, and Nutrient Availability**

Corn produces more residue than either soybeans or small grains. For example, a 150 bu/acre corn crop will produce about 8400 lbs residue/acre, whereas a 45 bu/acre soybean crop generates about 2500 lbs residue/acre, much of this being leaves which quickly decompose. A 60 bu/acre wheat crop will produce about 3600 lbs residue/acre. The large amount of residue from corn is an asset in building soil organic matter and protecting the soil from erosion. If current climate projections, i.e., more intense storms, hold true (Seeley, 2012), then the value of the residue becomes increasingly important. However, large amounts of residue can also pose challenges in creating a “good” seedbed, controlling pests, and recycling nutrients. The high level of corn residue is a concern for wheat because it acts as a host for the fungi *Fusarium graminearum*, which causes wheat head scab. For this reason, it is not a good idea to follow a corn crop with a wheat crop.

Soybeans tend to tolerate high-residue situations better than many other crops. The persistence of corn residue may slow nutrient recycling and the release of N from decaying stover. Following corn with a legume crop such as soybeans can be used to overcome this problem.

The use of cover crops before and following corn is a topic that needs additional investigation. Research at the SDSU Southeast Research Farm suggests that corn yields are higher following fall-planted, cool-season broadleaves (brassicas and legumes such as radish and peas) than grass-dominated cover crops (Sexton et al., 2012, 2014). Benefits of cover crops on corn yields is attributed to improved nutrient recycling and increased plant diversity (Sexton et al., 2009).

### **Impacts on Yield in a Corn-Soybean System**

Studies in South Dakota, Minnesota, Wisconsin, and Nebraska have reported a 10% to 22% yield benefit for corn grown in rotation with soybeans versus a continuous corn cropping pattern (Porter et al., 1997; Reidell et al., 2009; Stanger and Lauer, 2008; Wilhelm and Wortmann, 2004) (Fig. 9.2). Similar results were observed for soybeans where there was an 8% to 10% yield advantage when grown in rotation with corn rather than a continuous soybean rotation (Porter et al., 1997; Pederson and Lauer, 2004; Wilhelm and Wortmann, 2004). The rotational effect is attributed to many factors including enhanced root growth (Nickel et al., 1995).

Crop rotations can impact profitability. A 15-year Wisconsin study compared the corn-soybean rotation with continuous corn and rotations that contained oats and alfalfa (Stanger et al., 2008). This study reported that the corn-soybean rotation was more profitable than continuous corn and rotations that include oats and alfalfa.

While a corn-soybean rotation has been shown to be superior to continuous corn, it is still not a very diverse system. In many fields, there is a corn yield decrease of about 5% to 15% for second-year corn relative to first-year corn. The greatest yield reductions are typically measured between first- and second-year corn but can also be high when weather is unfavorable. Yield reductions generally stabilize after the third-year corn. Soybeans have similar responses and generally yield 5% to 8% more when following two or more years of corn. Crookston et al. (1991) conducted a 9-year study looking at corn and soybean yields in




Figure 9.2 Average corn yields, at the end of six years, in three rotational sequences. (Katsvairo and Cox, 2000) Even after just two cycles, yield differences were apparent. Data shown are averages across three tillage regimes from the final year of the study. These plots received 145 lbs/acre of N.

southwestern Minnesota. They concluded that “a superior cropping sequence ... would include at least three crops and possibly more.” Additional benefits from diverse rotations include reduced development of pest resistance, improved ability to manage variable weather conditions, and increased economic diversification.

### Rotations and Water Use

Rotations can be used to improve water management. For example, rotations provide protection from summer droughts by distributing the critical water-use periods across the growing season. Research conducted by the author shows that corn, wheat, and soybeans have different critical periods for water stress. Wheat partially avoids drought-stress by flowering and completing its lifecycle earlier in the growing season than either corn or soybeans (Fig. 9.3). Soybean flowering is spread over several weeks so that it can better avoid the effects of drought. The corn crop, on the other hand, flowers and sets seed at one point in time and does this during the warmest part of the year, when evaporative demand (water use) is at its peak. High temperatures and drought stress can reduce corn kernel set by decreasing pollen viability and delaying silking. By seeding hybrids of different maturities, the length of the pollination period for the farm can be expanded.




Figure 9.3 Estimated crop water used by spring wheat, corn, and soybeans grown over a season at Huron, S.D. Based on data from the University of Minnesota Extension, [www.extension.umn.edu](http://www.extension.umn.edu), and South Dakota climate archives, [www.climate.sdstate.edu](http://www.climate.sdstate.edu). Note: Drought stress is reduced by shifting wheat water use earlier in the season.

A worksheet for calculating agricultural intensity for different rotations is available at the South Dakota Lakes website. This calculator can be used to determine water harvesting from the different crops in a rotation. Along with water-use timing, crop rooting depth should be considered. Crops with deep extensive root systems that grow late into the season (e.g., sunflower and alfalfa) are likely to leave less reserve moisture than shallower-rooted, earlier maturing crops (e.g., peas, flax, and lentils). Cropping more frequently with high water-use crops increases the cropping system intensity. Barley, winter wheat, field peas, and canola are low water-use crops, whereas corn, soybean, and alfalfa are high water-use crops.

### Crop Diversity

When considering diversity, crop rotations can increase diversity and reduce problems with labor, equipment, disease, weeds, and insects. Diversity assessments should consider the type of plant. In South Dakota, commonly grown crops can be classified as:

1. Cool-season grass: spring wheat, winter wheat, barley, durum wheat, oat, and winter rye.
2. Warm-season grass: corn, sorghum, sudangrass, and millet.
3. Warm- and cool-season broadleaf plants such as field pea, lentil, canola, mustard, crambe, flax, safflower, chickpea, sugar beet, sunflower, dry edible bean, soybean, and alfalfa.

When selecting a crop rotation it is important to avoid potential conflicts between the seeding and harvest times of different crops (i.e., trying to seed one crop when harvesting another, or harvesting more than one crop at a time).

### Rules of Thumb for Increasing Diversity in Semi-arid Regions

1. Use soil survey information to evaluate soil water storage. Determine the appropriate cropping intensity based on this information.
2. Manage crop residues to facilitate soil water storage.
3. Manage crop nutrients to optimize yields while minimizing competition with weeds.
4. Utilize legume crops and animal manure to increase energy efficiency and improve soil quality.

5. Adopt techniques that minimize wind and water erosion.
6. Anticipate equipment and/or labor requirements for growing new crops.
7. Use cover crops to increase crop rotation intensity and diversity.
8. Consider a perennial crop, such as grass or alfalfa. They provide excellent weed suppression in a rotation, particularly if the crop following perennial plant is planted with minimal soil disturbance.
9. Consider the marketability of the commodity prior to planting a crop.
10. Avoid using crops with the same pests after each other. For example, soybeans should not follow field peas.

### **Importance of Linking Tillage and Crop Rotations**

Crop rotation and tillage should be considered simultaneously. Designing appropriate crop rotations is a mix of art and science. For any given situation, there will be a range of rotations that are appropriate. Within this range, there are rotations and tillage practices that reduce or increase risks. Additional information on tillage systems is available in Chapter 11.

### **References and Additional Information**

- Anderson, R.L. 2008. Diversity and no-till: keys for pest management in the U.S. Great Plains. *Weed Sci.* 56:141-145.
- Beck, D.L. 2003. Profitable No-till Systems Designed for Producers in the North American Great Plains and Prairies. Conference Proceedings, 2003 Sixth Annual Northwest Direct Seed Cropping Systems Conference. Pasco, WA.
- Clay, D.E., S.A. Clay, C.G. Carlson, and S. Murrell. 2012. Mathematics and Science for Agronomists and Soil Scientists. International Plant Nutrition Institute.
- Crookston, R.K., J.E. Kurle, P.J. Copeland, J.H. Ford, and W.E. Lueschen. 1991. Rotational cropping sequence affects yield of corn and soybean. *Agron. J.* 83: 108-113.
- Katsvairo, T.W., and W.J. Cox. 2000. Tillage X rotation X management interactions in corn. *Agron. J.* 92: 493-500.
- Nickel, S.E., R.K. Crookston, and M.P. Russelle. 1995. Root growth and distribution are affected by corn-soybean cropping sequence. *Agron. J.* 87: 895-902.
- Pedersen, P., and J.G. Lauer. 2004. Soybean growth and development response to rotation sequence and tillage system. *Agron. J.* 96: 1005-1012.
- Porter, P.M., J.G. Lauer, W.E. Lueschen, J.H. Ford, T.R. Hoverstad, E.S. Oplinger, and R.K. Crookston. 1997. Environment affects corn and soybean rotation effect. *Agron. J.* 89: 441-448.
- Riedell, W.E., J.L. Pikul, Jr., A.A. Jaradat, and T.E. Schumacher. 2009. Crop rotation and nitrogen input effects on soil fertility, maize mineral nutrition, yield, and seed composition. *Agron. J.* 101: 870-879.
- Seeley, M.W., 2012. Climate trends and issues in the North-Central region. Carbon, Energy, and Climate Conference Proceedings. USDA-SARE and Michigan State University, Kellogg Biological Station. Hickory Corner, MI.
- Sexton, P., R. Berg, and D. Beck. 2012. Effect of cover crops on corn N requirements in a drought year. South Dakota State University Agricultural Experiment Station. Southeast Research Farm Annual Progress Report 2012. Pp. 21-25.
- Sexton, P., D. Johnson, and B. Rops. 2014. Evaluation of effects of cover crops on corn N requirements in 2014. South Dakota State University Agricultural Experiment Station. Southeast Research Farm Annual Progress Report 2014. Pp. 30-32.

- Sexton, P.J., G.A. Porter, C.B. Fitzgerald, and B.R. Hoskins. 2009. Observation of an apparent increase in P availability for potatoes following a green manure. In 2009 Agronomy abstracts. ASA. Madison, WI.
- Stanger, T.F., and J.G. Lauer. 2008. Corn grain yield response to crop production and nitrogen over 35 years. *Agron. J.* 100: 643-650.
- Stanger, T.F., J.G. Lauer, and J.P. Chavas. 2008. The profitability and risk of long-term cropping systems featuring different rotations and nitrogen rates. *Agron. J.* 100: 105-113.
- Thirgood, J.V. 1981. *Man and the Mediterranean Forest: A History of Resource Depletion*. Pp. 194. Academic Press, Boston, Massachusetts.
- Varvel, G.E., and W.W. Wilhelm. 2003. Soybean nitrogen contribution to corn and sorghum in western Corn Belt rotations. *Agron. J.* 95: 1220-1225.
- Wilhelm, W.W., and C.S. Wortmann. 2004. Tillage and rotation interactions for corn and soybean grain yield as affected by precipitation and air temperature. *Agron. J.* 96: 425-432.

## Acknowledgements

Support for this document was provided by South Dakota State University, SDSU Extension, and the South Dakota Corn Utilization Council.



**A G R O W I N G I N V E S T M E N T**

Sexton, P. 2016. Chapter 9: Crop Rotations Can Increase Corn Profitability and Reduce Pests. In Clay, D.E., C.G. Carlson, S.A. Clay, and E. Byamukama (eds). iGrow Corn: Best Management Practices. South Dakota State University.

*The preceding is presented for informational purposes only. SDSU does not endorse the services, methods or products described herein, and makes no representations or warranties of any kind regarding them.*

*In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.*

*Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotope, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.*

*To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [http://www.ascr.usda.gov/complaint\\_filing\\_cust.html](http://www.ascr.usda.gov/complaint_filing_cust.html) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by:*

(1) mail: U.S. Department of Agriculture  
Office of the Assistant Secretary for Civil Rights  
1400 Independence Avenue, SW  
Washington, D.C. 20250-9410;

(2) fax: (202) 690-7442; or

(3) email: [program.intake@usda.gov](mailto:program.intake@usda.gov).

*USDA is an equal opportunity provider, employer, and lender.*

SDSU Extension is an equal opportunity provider and employer in accordance with the nondiscrimination policies of South Dakota State University, the South Dakota Board of Regents and the United States Department of Agriculture