

## BEST MANAGEMENT PRACTICES

### Chapter: 20 Corn Nitrogen Timing



Christopher Graham (Christopher.Graham@sdstate.edu) and Anthony Bly (Anthony.Bly@sdstate.edu)

To optimize uptake of nitrogen (N) fertilizer efficiency and to minimize the adverse impact of N on the environment, we recommend that N be applied at the right time, in the right form, at the right place, and in the right amount. This chapter specifically addresses applying N at the right time. A corn plant takes up a large percentage of its N between the V6 (~10 inch tall) and R1 (silking) growth stages. During this period, newer hybrids require as much as 8 lbs/day to maintain maximum production. When N is applied earlier than it is needed by the plant, it can be lost through a variety of mechanisms including leaching and denitrification.

**Table 20.1 Advantages and disadvantages of various fertilizer timing.**

N Timing	Advantage	Disadvantage
Fall	Equipment is available.	N loss off-field can be through several mechanisms.
	Weather permitting, time is typically not a constraint.	Applied prior to crop demand. Often has lower efficiency than spring and split applications.
Preplant spring	Reduced N losses relative to fall.	If excessive spring rainfall, N loss off-field can still be significant.
At planting pop-up and starter fertilizer	Applied with or near the seed.	Salts and ammonium/ammonia in the fertilizer can inhibit germination.
	For pop-up use, only low rates required.	Applied prior to crop demand.
Topdressed, applied to standing crop after emergence and when corn is < 6 inches	Reduces losses relative to fall or preplant spring.	Volatilization losses may be high when surface broadcast.
		Rainfall required to move N into the soil when surface broadcast.
		Leaching/denitrification losses may be high following spring rainfalls, but inhibitors may help.
Sidedressed, applied when corn is <12 inches tall	Applied when plant needs N.	Accounts for early spring rainfall, however, rainfall is required to move the N into the soil.
	Accounts for rainfall.	Can be delayed by excessive rainfall conditions.
	Improved N efficiency.	Time demands due to weed control requirements.
Combination of starter + sidedressed (Split)	Accounts for rainfall, can have high efficiency.	Time required to apply the sidedressed N.

The advantages and disadvantages associated with N fertilizer timing in corn production are summarized in Table 20.1. Corn producers need to consider weather, N fertilizer source, placement, and cultural practices such as tillage, pest, weed, and disease pressure. The ultimate goal of an N fertilization program is to supply N when it is most needed. While economic and logistic factors make fall N applications more convenient, the practice has risks that may not be worth the trade-offs. In years with a wet spring a significant amount of N may be lost, making spring and split N applications preferable. This chapter provides management guidelines for fall, spring, and split N applications.

### **N Fertilizer Options**

With the high cost of N fertilizer and an awareness of adverse effects of N on the environment, there is an increased interest in adopting techniques that improve N fertilizer efficiency. Recent research has strengthened the case that in humid or irrigated environments, a split application is more effective at meeting corn N demands than a single application applied either in the fall or early spring. However, in rainfed corn production systems, delaying the N application increases the risk that surface-applied N will not be incorporated and available for uptake from crop roots. When this happens, the yield loss can also be substantial.

#### ***Fall Broadcast Applications of Urea***

Broadcast-applied urea is most susceptible to environmental loss through volatilization. It is not recommended to fall apply N before the soil cools to less than 50°F or to sandy soil. Moreover, research from Montana found that application to snow-covered soil still maintained fairly high volatilization rates, particularly during periods of snowmelt. Additionally, application to soil with high pH, generally above 7.5, increases volatilization. Ammonia volatilization is reduced by incorporating the urea granular into the soil either by cultivation or rainfall.

#### ***Starter and Pop-up Fertilizers***

Pop-up fertilizers are placed with the seed at planting, whereas starter fertilizer is applied near the seed. Salts contained within the fertilizer can reduce seed germination. For pop-up fertilizers, avoid the use of urea. For corn production in South Dakota, di-ammonium phosphate (DAP) or mono-ammonium phosphate (MAP) are recommended as a pop-up fertilizer. Both of these products contain a small amount of N. Generally, the  $K_2O + N$  rate applied with the seed should not exceed 10 lbs/acre, however this rate is dependent on row spacing and soil texture. A calculator to determine pop-up fertilizer rates is available from the International Plant Nutrition Institute, [ipni.net/article/IPNI-3268](http://ipni.net/article/IPNI-3268).

Starter fertilizer is generally placed 2 inches to the side and 2 inches below the seed. By separating the fertilizer and seed, the risk of salt injury is reduced. However, this risk is not eliminated and it is recommended to apply less than 70 lbs of  $N + K_2O$  if the band is within 2 inches of the seed.

#### ***Split-N Applications***

In a split-N application system, N is applied at multiple times during the season. It can be applied in the fall, with the seed as a starter, in a band next to the seed (2 inches deep by 2 inches to the side of the seed row), or between the rows as a sidedressed application. One of the greatest strengths with the split-N approach is that it allows the grower to account for early season N losses and changes in the grain yield potential. When a pre-season nitrogen test shows adequate soil N, a producer may benefit economically by reducing preplant or starter N rates.

Research in South Dakota indicates that splitting N applications between preplant and V6 can increase corn grain yield over fall application of N. However, the amount of N to apply at V6 is dependent on the amount of nitrate-N contained in the soil. The pre-sidedress nitrate test (PSNT) is one tool that can be used to estimate the sidedress N application rate. For PSNT, randomly collect 16 to 24 soil cores from the surface 12 inches when the plants are between V3 to V5 (Magdoff et al., 1984). Sample collection and handling should follow good sampling protocols. Determine the N rate based on the  $NO_3^-$ -N concentration (Table 20.2).

**Table 20.2 Relationship between the amount of in-season soil test N in the surface 12 inches and the sidedressed N rate. (Modified from Blackmer et al., 1991; Reitsma et al., 2008)**

In-season soil test NO <sub>3</sub> -N	Sidedressed N
ppm	lbs N/acre
0-10	80-120
11-15	50-90
16-20	30-60
21-25	0-40
> 25	0

One of the problems with applying N at the V6 growth stage is that rainfall is needed to move the fertilizer into the soil where it can be absorbed by the roots. This problem can be avoided by injecting the N into the soil. However, if the in-season N application is delayed due to high rainfall or logistical issues, recent research from Missouri suggests that “rescue N” applications can be applied as late as tasseling. However, growers must take precautions to minimize leaf burn.

#### Summary of N Timing

Nitrogen is typically lost through volatilization, leaching and/or denitrification. A single, fall application of N represents a gamble on whether N will be available in late

June and early July when it is most needed by the crop. There are cases where it has been found acceptable (Bundy, 1986; Vetsch and Randall, 2004). However, fall applications on sandy soils are not recommended, and it is not recommended on other soils until temperatures decrease below 50°F.

The potential losses from early-applied N and the yield advantage from in-season application are well-defined. After accounting for N in the soil, a broader set of recommendations would include N application through a combination of applying some N (30-50 lbs/acre) early in the growing season, with the remainder being applied in the mid-vegetative stages. Protecting the N with urease and nitrification inhibitors can also prolong the time period when N is safe from loss. Where possible, adding N through irrigation water is an effective approach.

## References and Additional Information

- Bender, R.R., J.W. Haegele, M.L. Ruffo, and F.E. Below. 2013. Nutrient uptake, partitioning, and remobilization in modern, transgenic insect-protected maize hybrids. *Agron. J.* 105:161-170.
- Binford, G.D., A.M. Blackmer and M.E. Cerrato. 1992. Relationships between corn yields and soil nitrate in late Spring. *Agron J.* 84: 53-59.
- Blackmer, A.M., D. Pottker, M.E. Cerrato and J. Webb. 1989. Correlations between soil nitrate concentrations in late spring and corn yields in Iowa. *J. Prod. Ag.* 2:103-109.
- Blackmer, A.M., T.F. Morris, D.R. Keeney, R.D. Voss, and R. Killorn. 1991. Estimating nitrogen needs for corn by soil testing. PM1381. Iowa State University, Iowa CES, Ames.
- Bly A., J. Gerwing and R. Gelderman. 2004. Nitrogen application timing and rate influence on nitrate-nitrogen leaching and corn grain yield near Aurora SD during 2004. Soil/Water Research, South Dakota State University Agriculture Experiment Station TB099 (SOIL PR 04-30).
- Bundy, L.G. 1986. Review: Timing nitrogen applications to maximized fertilizer efficiency and crop response in conventional corn production. *J. Fert. Issues* 3:99-106.
- Cassman, K.G., A. Dobermann and D.T. Walters. 2002. Agroecosystems, nitrogen-use efficiency and nitrogen management. *Ambio* 31:132-140.
- Gelderman, R., J. Gerwing and A. Bly. 2006. Nitrogen management in a corn/soybean rotation. Soil/Water Research, South Dakota State University Agriculture Experiment Station TB099 (SOIL PR 06-4).
- Magdoff, F.R., D. Ross and J. Amadon. 1984. A soil test for nitrogen availability to corn. *Soil Sci. Soc. Am. J.* 48:1031-1304
- Reitsma, K.D., G. Gelderman. P. Skiles, K. Alverson, J. Hemenway, H.J. Woodard, T.E. Schumacher, D. Malo, and D.E. Clay. 2008. Nitrogen best management practices for corn in South Dakota. South Dakota Extension GS941.
- Roberts, T.L. 2007. Right product, right rate, right time, and right placement... foundation of best management practices for fertilizers. International Plant Nutrition Institute available at [http://www.ipni.net/publication/bettercrops.nsf/0/91607AF3210A609F852579800080C01C/\\$FILE/Better%20Crops%202007-4%20p14.pdf](http://www.ipni.net/publication/bettercrops.nsf/0/91607AF3210A609F852579800080C01C/$FILE/Better%20Crops%202007-4%20p14.pdf) (accessed September 4, 2015).
- Sogbedji, J.M., H.M. van Es, C.L. Yang, L.D. Geohring, and F.R. Magdoff. 2000. Nitrate leaching and N budget as affected by maize N fertilizer rate and soil type. *J. Environ. Qual.* 29:1813-1820.
- Vetsch, J.A. and G.W. Randall. 2004. Corn production as affected by nitrogen application timing and tillage. *Agron. J.* 96:502-509.

## Acknowledgements

Support for this document was provided by South Dakota State University, SDSU Extension, and the South Dakota Corn Utilization Council.



**A G R O W I N G I N V E S T M E N T**

Graham, C., and A. Bly. 2016. Chapter 20: Corn Nitrogen Timing. In Clay, D.E., C.G. Carlson, S.A. Clay, and E. Byamukama (eds). *iGrow Corn: Best Management Practices*. South Dakota State University.

*The preceding is presented for informational purposes only. SDSU does not endorse the services, methods or products described herein, and makes no representations or warranties of any kind regarding them.*

*In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.*

*Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.*

*To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [http://www.ascr.usda.gov/complaint\\_filing\\_cust.html](http://www.ascr.usda.gov/complaint_filing_cust.html) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by:*

(1) mail: U.S. Department of Agriculture  
Office of the Assistant Secretary for Civil Rights  
1400 Independence Avenue, SW  
Washington, D.C. 20250-9410;

(2) fax: (202) 690-7442; or

(3) email: [program.intake@usda.gov](mailto:program.intake@usda.gov).

*USDA is an equal opportunity provider, employer, and lender.*

SDSU Extension is an equal opportunity provider and employer in accordance with the nondiscrimination policies of South Dakota State University, the South Dakota Board of Regents and the United States Department of Agriculture