

Chapter: 48 Corn Foliar Fungicides



Emmanuel Byamukama (Emmanuel.Byamukama@sdstate.edu) and
Connie Strunk (Connie.Strunk@sdstate.edu)

Foliar diseases can lead to premature leaf senescence, and predispose stalks to rotting, poor grain quality, and reduced yields. Common fungal diseases found on corn include common rust, northern corn leaf blight, gray leaf spot, eye spot, anthracnose leaf blight, and Physoderma brown spot. Management of foliar diseases involves managing the surface residue (through rotation or tillage), selecting resistant hybrids, and performing in-season fungicide application. Corn residue on the surface of the soil can increase certain foliar disease problems, such as gray leaf spot and northern corn leaf blight. Although the severity of these diseases varies from year to year, application of foliar fungicides may provide effective control in years of high disease pressure. The purpose of this chapter is to provide guidance on the use of fungicides.



Figure 48.1 Northern corn leaf blight symptoms on corn. (Courtesy of authors)

Introduction

Fungicides are an effective in-season management tool for fungal leaf diseases, and sometimes can reduce chances of stalk rot development. A number of fungicide products that are effective against fungal pathogens on corn are available for use. However, some are more effective on certain pathogens than others. Most of the fungicides available are preventive in nature and stop the fungus from infecting or advancing within the plant. Therefore, timing of a fungicide treatment is critical. If fungicides are applied when the severity is already high, the benefit will be limited.

When deciding whether to apply a foliar fungicide, consider the following:

- The level of disease. Is there a significant amount of disease showing up on the leaves below the ear leaf?
- The current weather. For example, has it been warm and humid? Does the forecast predict continued hot conditions? If yes, disease severity may worsen, so application is advised. If no, disease outbreaks may not reach a critical stage and scouting should continue until corn has passed dent growth stage.
- The corn growth stage. How far along is the corn? If corn is at R5 (dent), diseases most likely will not

influence yield or will be minimal.

- Susceptibility of hybrid(s). For example, most the hybrids have moderate resistance to common rust and therefore, no treatment may be needed.
- Potential yield. If yield is predicted to be low (due to moisture stress, or poor fertility), chances of an economic gain due to fungicide treatment will be low.
- Grain price. When prices are high, it takes only a few bushels to pay for the cost of applying a fungicide (Table 48.1).

Once a fungicide treatment is deemed to be necessary, growers should ensure the sprayer is calibrated to deliver the recommended rate (as per the fungicide label), and that weather conditions are not too windy (> 10 mph) or too hot.

Table 48.1 The number of corn bushels needed to break even for the cost of fungicide and its application.

Price of corn (\$)	Application cost (\$)						
	12.0	15.0	20.0	25.0	30.0	40.0	50.0
1.5	8.0	10.0	13.3	16.7	20.0	26.7	33.3
2.0	6.0	7.5	10.0	12.5	15.0	20.0	25.0
3.0	4.0	5.0	6.7	8.3	10.0	13.3	16.7
4.0	3.0	3.8	5.0	6.3	7.5	10.0	12.5
5.0	2.4	3.0	4.0	5.0	6.0	8.0	10.0
6.0	2.0	2.5	3.3	4.2	5.0	6.7	8.3
7.0	1.7	2.1	2.9	3.6	4.3	5.7	7.1
8.0	1.5	1.9	2.5	3.1	3.8	5.0	6.3
9.0	1.3	1.7	2.2	2.8	3.3	4.4	5.6

Proactive Fungicide Treatments

Economic Benefit

Several research studies have shown that when a fungicide is applied in the absence of disease or very low disease severity, the probability of increasing yield to pay for the treatment decreases significantly (Byamukama et al, 2013; Wise and Mueller, 2011; Pierce et al, 2011). For example, Mueller and Wise (2011) analyzed data from 613 treatment comparisons of strobilurin-treated and nontreated plots over a 10-year period in the Corn Belt region. The fungicides were applied between V14 and R5 (dent) with a majority of treatments being applied between tasseling (VT) and R2 (blister). They reported that when disease severity was less than 5% on the ear leaf at the end grain fill period, the fungicide treatment increased yields 1.5 bu/acre, and when the disease severity was > 5% the yield gain averaged 9.6 bu/acre. These results suggest that there may be some benefit from proactive fungicide applications. The yield enhancement has been linked to improved crop health (stays green longer) and reduced fungal populations. However, these benefits must be balanced against the long-term risk of the fungal pathogens developing resistance. Therefore, to avoid problems associated with unnecessary application of fungicides (such as resistance development, added expenses), growers should always scout to determine the need for a fungicide application.

Fungicide Efficacy for Control of Corn Diseases - 2016

The South Dakota State University Plant Pathology Extension is a member of the Corn Disease Working Group (CDWG) and has participated in the fungicide efficacy trials. The group has developed the following information on fungicide efficacy for management of major corn diseases in the United States. Efficacy ratings for each fungicide listed in the table were determined by committee members field-testing the materials over multiple years and at multiple locations. Efficacy ratings are based upon level of disease control achieved by product and are not necessarily reflective of yield increases obtained from product application. Efficacy depends upon proper application timing, rate, and application method to achieve optimum effectiveness of the fungicide as determined by labeled instructions and overall level of disease in the field at the time of application. Differences in efficacy among fungicide products were determined by direct comparisons among products in field tests and are based on a single application of the labeled rate as listed in Table 48.2.

Table 48.2 Systemic fungicides available that have been tested over multiple years and at multiple locations. The table is not intended to be a list of all labeled products¹. Efficacy categories: NR=Not Recommended; P=Poor; F=Fair; G=Good; VG=Very Good; E=Excellent; NL=Not Labeled for use against this disease; U=Unknown efficacy or insufficient data to rank product. This table is a joint publication by the Corn Diseases Working Group, coordinated by Dr. Kiersten Wise at Purdue University. Other recommendations can be found at extension.sdstate.edu.

Fungicide (s) Class	Active ingredient (%)	Product/Trade name	Rate/a (fl oz)	Anthraco nose leaf blight	Common rust	Eye spot	Gray leaf spot	Northern leaf blight	Southern rust	Harvest Restriction ²
QoI Strobilurins Group 11	Azoxystrobin 22.9%	Quadris 2.08 SC Multiple Generics	6.0-15.5	VG	E	VG	E	G	G	7 days
	Pyraclostrobin 23.6%	Headline 2.09 EC/SC	6.0-12.0	VG	E	E	E	VG	VG	7 days
	Picoxystrobin	Aproach 2.08 SC	3.0-12.0	VG	VG-E	VG	F-VG	VG	G	7 days
DMI Triazoles Group 3	Propiconazole 41.8%	Tilt 3.6 EC Multiple Generics	2.0-4.0	NL	VG	E	G	G	F-G	30 days
	Prothioconazole 41.0%	Proline 480 SC	5.7	U	VG	E	U	VG	G	14 days
	Tebuconazole 38.7%	Folicur 3.6 F Multiple Generics	4.0-6.0	NL	U	NL	U	VG	F-G	36 days
	Tetraconazole 20.5%	Domark 230 ME	4.0-6.0	U	U	U	E	U	G	R3 (milk)
Mixed modes of action	Azoxystrobin 13.5%	Quilt Xcel 2.2 SE	10.5- 14.0	VG	VG-E	VG-E	E	VG	VG	30 days
	Propiconazole 11.7%									
	Bensovindiflupyr 10.27%	Trivapro A 0.83 + Trivapro B 2.2 SE	A = 4.0 B = 10.5	U	U	U	E	VG	E	7 days (A) 30 days (B)
	Azoxystrobin 13.5%									
	Propiconazole 11.7%									
	Cyproconazole 7.17%	Aproach Prima 2.34 SC	3.4-6.8	U	U	U	E	VG	G-VG	30 days
	Picoxystrobin 17.94%									
Flutriafol 19.3%	Fortix 3.22 SC Preemptor 3.22 SC	4.0-6.0	U	U	U	E	VG-E	VG	R4 (dough)	
Fluoxastrobin 14.84%										
Pyraclostrobin 28.58%	Priaxor 4.17 SC	4.0-8.0	U	VG	U	VG	U	G	21 days	
Fluxapyroxad 14.33%										

Fungicide (s) Class	Active ingredient (%)	Product/Trade name	Rate/a (fl oz)	Anthraco nose leaf blight	Common rust	Eye spot	Gray leaf spot	Northern leaf blight	Southern rust	Harvest Restriction ²
Mixed modes of action	Pyraclostrobin 13.6%	Headline AMP 1.68 SC	10.0–14.4	U	E	E	E	VG	G-VG	20 days
	Metconazole 5.1%									
	Trifloxystrobin 32.3%	Stratego YLD 4.18 SC	4.0–5.0	VG	E	VG	E	VG	G-VG	14 days
Prothioconazole 10.8%										
	Tetraconazole 7.48%	Affiance 1.5 SC	10.0–14.0	U	U	U	U	U	G	7 days
	Azoxystrobin 9.35%									

¹Additional fungicides are labeled for disease on corn, including contact fungicides such as chlorothalonil. Certain fungicides may be available for diseases not listed in the table, including Gibberella and Fusarium ear rot. Applications of Proline 480 SC for use on ear rots requires a FIFRA Section 2(ee) and is only approved for use in Illinois, Indiana, Iowa, Louisiana, Maryland, Michigan, Mississippi, North Dakota, Ohio, Pennsylvania, and Virginia.

²Harvest restrictions are listed for field corn harvested for grain. Restrictions may vary for other types of corn (sweet, seed or popcorn, etc.), and corn for other uses such as forage or fodder.

Many products have specific use restrictions about the amount of active ingredient that can be applied within a period of time or the amount of sequential applications that can occur. Please read and follow all specific use restrictions prior to fungicide use. This information is provided only as a guide. It is the responsibility of the pesticide applicator by law to read and follow all current label directions. Reference to products in this publication is not intended to be an endorsement to the exclusion of others that may be similar. Persons using such products assume responsibility for their use in accordance with current directions of the manufacturer. Members or participants in the CDWG assume no liability resulting from the use of these products.

Fungicide Resistance

Fungicide resistance is when a fungicide that used to control a given fungal pathogen, no longer offers any protection against the same fungus. Several factors are responsible for fungicide resistance including:

- The fungicide provides a selection process for pathogens that are resistant or tolerant to the treatment. Practices that increase the risk of fungicide resistance include:
 - Multiple applications of fungicides with same mode of action.
 - Reducing the application rate or using off-label products.
 - Multiple applications of the same mode of action fungicides within a single year.
- High genetic variability within the pathogens.
 - High variability suggests that some pathogens will have inherent tolerance to fungicide.
- High reproduction capacity of the pathogen.
 - Pathogens, which reproduce quickly (e.g., rusts), are likely to have increased diversity and therefore likely to be selected for when fungicides of similar modes of action are applied to the same area in a season.

To avoid fungicide resistance, growers should monitor the performance of the fungicides they use. One way to do this is to leave a strip of a nontreated area (one pass), where the yield and disease severity from the treated and nontreated zones can be compared. If the two areas have comparable disease severity, this would mean that the fungicide had minimal impact on the disease that year and this could probably be due to resistance development. In this case, samples of diseased leaves in the treated area and untreated area should be collected and sent to the SDSU Plant Diagnostic Clinic for fungicide sensitivity testing. Proper disease identification and appropriate fungicide selection is crucial for effective use of fungicides.

The risk of developing fungicide resistance can be reduced by:

- Rotating between different classes of fungicide within a season and also between seasons.
- Scouting to determine the need for a fungicide and avoid applying fungicide when it is not necessary or when it is too late (severe symptom on ear leaf and higher).
- Using a mixture of fungicide classes. Luckily, several fungicide products are “broad spectrum.”
- Practicing integrated disease management to reduce disease pressure.
- Following the label directions to determine the rates, growth stage of the crop, compatibility with other pesticides, and safety information.

Fungicide Classes

Fungicides are classified into groups depending on their mode of action. For instance, some fungicides interfere with fungal protein synthesis, while others interfere with respiration, etc. The Fungicide Resistance Action Committee (FRAC) is an international committee that is responsible for fungicide-resistance monitoring. The panel has developed FRAC codes that classify fungicides into classes with the same mode of action (Fig. 48.2). Fungicide labels contain the FRAC code and fungicides with the same FRAC code belong to the same class. When rotating fungicides, growers should ensure that rotation is made between different FRAC codes. Fungicide resistance has not been reported for corn pathogens in South Dakota.

Host Resistance in Management of Plant Diseases

Cultivar selection is a critical step in integrated pest management (IPM). Prior to the use of synthetic chemicals, farmers chose and saved seed from the best yielding and healthiest plants (e.g., bigger corn ears) for the next growing season. Today, planting a carefully selected corn hybrid may be the most important management decision to get maximum yield. Corn hybrids are developed to suit different needs: maturity, resistance to pests and diseases, plant characteristics (plant height, seed color, stalk strength, etc.).

Host resistance/tolerance, when available, is the first line of defense in plant disease management. Disease resistance genes have been bred into hybrids through conventional breeding or genetic engineering. Evolutionarily, plants and pathogens have co-existed together. When plants are attacked by a pathogen, the

plants have evolved and developed a resistance gene against this pathogen. Over time, this pathogen also evolves to overcome the resistance gene.

Growers should keep good records on the performance of the hybrids grown to aid in their decision-making process. Monitoring performance of hybrids planted may also help indicate development or change in a pathogen race allowing the pathogen to overcome the resistance gene in the hybrid.

Unlike other traits, like glyphosate resistance, Bt, and other GMO traits, disease-resistance traits, to date, do not add to the cost of seed. Yet host resistance is an effective, sustainable, and affordable plant disease management practice. Several seed companies provide disease ratings for their hybrids making it easier to choose optimum characteristics for the growing conditions. When selecting a hybrid, growers should consider the history of diseases in their fields and cropping practices (such as corn on corn or no-till). For instance, corn on corn under irrigation is likely to have Goss's wilt develop; therefore, a grower in this case would want to plant a Goss's wilt resistant/tolerant corn hybrid.




Figure 48.2 Example of fungicide labels displaying the FRAC code (circled in blue). Prosaro has FRAC code 3 while Aproach has code 11. These two fungicides belong to two different groups and therefore can be rotated to prevent/manage resistance.

References and Additional Information

Websites

<http://www.sdaba.org/agronomyconference/pdfs/FungicidesOsborne.pdf> (accessed 6 June 2016).

Paper and Manuals

Byamukama, E., Abendroth, L. J., Elmore, R. W., and Robertson, A. E. 2013. Quantifying the effect of pyraclostrobin on grain-fill period and kernel dry matter accumulation in maize. Online. Plant Health Progress doi:10.1094/PHP-2013-1024-02-RS.

Clay, D.E., S.A. Clay, and K. Reitsma. 2009. South Dakota Corn Best Management Practices. SDSU, Brookings, SD.

Mueller, D., and R. Pope. 2009. Field Guide: A Reference for Identifying Diseases of diseases, insects pests of corn. Iowa State University of Science and Technology. Available at: <http://www.agronext.iastate.edu/corn/docs/corn-field-guide.pdf>. (accessed 6 June 2016).

Mueller D., Wise, K., Dufault, N., Bradley, C., and Chilvers, M. 2013. Fungicides for field crops. APS Press. 120p.

Paul, P. A., L.V. Madden, C.A. Bradley, A.E. Robertson, G.P. Munkvold, G. Shaner, K. Wise, D.K. Malvick, T.W. Allen, A. Grybauskas, P. Vincelli, and P. Esker, P. 2011. Meta-analysis of yield response of hybrid field corn to foliar fungicides in the U.S. Corn Belt. *Phytopathology* 101: 1122-1132.

Wise, K., and Mueller, D. 2011. Are fungicides no longer just for fungi? An analysis of foliar fungicide use in maize. Online. APSnet Features

Acknowledgements

Support for this document was provided by South Dakota State University, SDSU Extension, South Dakota Corn Utilization Council, and the USDA-AFRI-IPM.



A G R O W I N G I N V E S T M E N T

Byamukama, E., and C. Strunk. 2016. Chapter 48: Corn Foliar Fungicides. In Clay, D.E., C.G. Carlson, S.A. Clay, and E. Byamukama (eds). *iGrow Corn: Best Management Practices*. South Dakota State University.

The preceding is presented for informational purposes only. SDSU does not endorse the services, methods or products described herein, and makes no representations or warranties of any kind regarding them.

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at http://www.ascr.usda.gov/complaint_filing_cust.html and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by:

(1) mail: U.S. Department of Agriculture
Office of the Assistant Secretary for Civil Rights
1400 Independence Avenue, SW
Washington, D.C. 20250-9410;

(2) fax: (202) 690-7442; or

(3) email: program.intake@usda.gov.

USDA is an equal opportunity provider, employer, and lender.

SDSU Extension is an equal opportunity provider and employer in accordance with the nondiscrimination policies of South Dakota State University, the South Dakota Board of Regents and the United States Department of Agriculture