

BEST MANAGEMENT PRACTICES

Appendix A: Corn Planting Guide



Obtaining maximum profit from a corn crop depends on the timely planting of an appropriate hybrid, at the proper depth, with a planter that evenly spaces the seed. The success of a corn crop is dependent on equipment maintenance, seedbed preparation, the development of a sound fertility and pest management program, and planting the seed. Early planting is best, but temperatures should be warm enough to assure quick germination and emergence, and late enough to avoid hard frosts. Planting opportunity windows can be narrow due to spring rains or a late warm-up. Time spent in the off-season maintaining equipment and planning tentative season-long schedules can increase efficiency in the spring (when time is limited). This section discusses planter maintenance, planting date, replanting considerations, seeding rate, and planting depth.

Planter Maintenance and Preparation

A corn planter is a piece of precision equipment, with each component working together to place the seed in the ground at a uniform depth and with a uniform distance between seeds. Research has shown that the uniform spacing of seed can increase yields up to 20 bu/acre (Doerge and Hall 2000). Although they are conducted too late to correct an in-season problem, stand counts and population surveys can be useful for determining if a planter should be calibrated prior to the next use. Growing conditions should also be evaluated as poor seed quality, or problems such as soil crusting, areas that are too wet or too dry, or cold soil temperatures for extended periods, may be responsible for non-uniform stands. Potential yield losses due to uneven stands can be estimated (Carlson et al. 2000). If planter calibration is necessary, always follow the manufacturer's instructions for calibrating seed metering equipment. Assistance is available from local Extension educators, crop consultants, or seed dealers.

Table A.1. Planter Maintenance Checklist

- ✓ Review owner's manual.
- ✓ Replace worn parts.
- ✓ Calibrate seed meters.
- ✓ Calibrate planter fertilizer and pesticide applicators.
- ✓ Check down pressure springs.
- ✓ Maintain even and recommended tire pressure.
- ✓ Lubricate bearings and other moving parts.

During planting, it is important to place seed at the proper depth and ensure that the walls of the furrow are not smeared by the opener. Down pressure tension should be adjusted if seed is not placed at the desired depth (1½ to 2") (see "Depth and Planting Options" section at the end of this chapter). Closers or packing wheels should apply enough pressure for good seed-to-soil contact; too much pressure

Table A.2. Suggested and historical dent corn planting dates in South Dakota by region

Approximate planting dates by reporting region						South Dakota Reporting Region
Suggested planting dates*			Historical acres planted, 1970 – 1994**			
Earliest	Latest	Desired range	10%	50%	90%	
May 4	Jun 5	May 12 -26	May 10	May 26	Jun 9	Northwest North Central Northeast
5	5	May 10 - 24	9	20	5	
6	5	May 10 - 24	6	18	4	
Apr 29	Jun 8	May 12 - 24	May 12	May 25	Jun 10	West Central Central East Central
May 3	5	May 6 - 26	9	20	5	
6	5	May 6 - 26	4	16	3	
May 4	Jun 3	May 7 - 24	May 7	May 20	Jun 2	Southwest South Central Southeast
Apr 29	8	May 3 - 17	10	22	7	
27	10	May 1 - 15	6	15	2	

* Dates are best estimates obtained from historical and research data within a reporting region

** Adapted from National Agricultural Statistics Service (NASS) – South Dakota Field Office

will compact the seedbed. Adjust down pressure tension in consideration of soil moisture and residue conditions.

As no-till and reduced-till systems become increasingly popular, the planter takes on the additional task of manipulating soil and crop residue. Hence, there are more parts to wear out and maintain. Implements that manage residue on the planter are critical in no-till and other high-residue systems, as crop residue can interfere with openers and closures.

Planting Dates

The spring planting window generally ranges from late April to mid-June (table A.2). Historically, 10% of the corn acres in South Dakota are seeded by mid-May and continuing to mid-June. Seed germination depends on soil moisture and temperature. Care should be taken to avoid tillage and planting operations when soil is wet. Yields may or may not be reduced due to delayed planting. However, due to problems associated with compaction, “mudding” the seed in will reduce the yield both of the current years’ crop and of those crops grown in the future.

As a general rule, corn should not be planted until the soil temperature (measured at 2” between 7 and 8 a.m.) approaches 50°F. In cold soil conditions (below 50°F), seeds will readily absorb water but will not initiate root or shoot growth; this leads to seed rots and poor emergence. If circumstances force planting before soil temperatures reach 50°F, it is recommended to consult with a reputable seed dealer or agronomist to select

Table A.3. Yield response of corn to planting date

Relative Maturity (MN Rating)	-----Average planting date-----					Daily yield loss from May 7 (bu/Acre)
	April 17	April 27	May 7	May 17	May 27	
	-----Average yield (bu/Acre)-----					(bu/Acre)
101 – 103 d. (early)	130	132	131	132	119	0.06
112 – 118 d. (late)	143	145	141	131	109	1.6
Average	137	139	136	131	114	1.1

*No data for 1995 or 2000

Yield data collected from 1986 to 2001 (14 yrs*)

Southeast South Dakota Experiment Station, Beresford SD

(Berg et al. 2001)

Table A.4. Estimated accumulated GDUs required for corn

Growth Stage	RM* – 80 days (Early)	RM* – 95 days (Mid)	RM* – 110 days (Late)
	-----GDUs-----		
Emergence	110	110	110
R1 (silking)	1100	1250	1400
R6 (maturity)	1900	2200	2500

* Relative Maturity (RM) of hybrid in days

an appropriate hybrid (one where the seed has been treated with a fungicide).

Delayed Planting or Replanting Considerations

Delayed planting reduces the number of growing degree units (GDU) accumulated during the season, hindering the crop from maturing before the first fall killing frost (see Appendix B). Corn killed by frost before maturity may not have completely filled kernels and has a slower dry-down rate, which can lead to excessive drying costs. If planting is delayed, late-maturing hybrids can lose up to 1.1 bu/Acre* per day compared to earlier-maturing hybrids that can be planted later in the season without realizing a loss (table A.3). Often, the trade-off is that earlier hybrids may inherently have a lower yield potential.

The number of GDUs that a hybrid needs to reach physiological maturity is related to maturity ratings (table A.4.). Since GDUs are based on temperature, the amount of GDUs accumulated in the spring and fall are less than during the peak summer months. Available GDUs decline with later planting dates. However, corn will usually emerge quicker if soil temperatures are warmer. If planting is delayed, an earlier maturing hybrid should be considered.

A rule of thumb is to plant 20% of fields with a full season hybrid, 60% with a mid-season hybrid, and the remaining 20% with a short-season hybrid (“20-60-20 rule”). If planting is delayed, growers are urged to consult their seed dealer to determine if an earlier-maturing hybrid is warranted.

Seeding Rates

The optimal population for an area is influenced by available water, nutrients, and overall soil productivity. Even within a field, optimal populations may vary by soil type or landscape position. Low populations can lead to increased weed pressure (from lack of competition), whereas higher plant populations increase seed investment with little return. Achieving an optimal population throughout the field gives corn a competitive edge over weeds and can optimize grain dry-down time in the fall.

Optimal corn populations vary from 24,000 to 32,000 plants per acre. Higher-productivity soils with sufficient drainage and available water can support higher populations. Data in table 3.5 provides a guide for selecting optimal population rates.

Some overall recommendations for seeding rate include:

- Increase populations by $\approx 10\%$ for silage crops.
- Set seeding rates higher than target population to account for less than 100% germination and seedling mortality.
- Increase seeding rate by ≈ 2000 seeds/acre in no-till systems.
- Increase seeding rate by ≈ 2000 to 3000 seeds/acre in irrigated fields.

Depth and Planting Operations

Depending on field conditions at the time of planting, depth can vary from 1- $\frac{1}{2}$ to 3 inches. Under optimal conditions, seed is commonly placed 1- $\frac{1}{2}$ to 2 inches below the soil surface. In dry conditions, it may be advantageous to plant deeper (2 to 3”) to place the seed into a higher-moisture area. If soil is very dry and rain is not expected, seed may be placed up to 3 inches deep. Planting deeper than 3 inches is not recommended because placing the seed too deep (>3”) will not allow emergence (as the coleoptile cannot elongate enough to bring the leafy parts above ground). Although soil conditions may be dry, consider the probability of rain in the near future. Rain can seal the surface of the soil, making it difficult for the developing plant to emerge. Shallower depths (<2”) should be targeted if rain is likely.

Crop residue can affect seeding date (as soils warm more slowly in high residue systems). Seed can be left on the surface when seed openers “ride-up” over residue. When seeding into areas with heavy residue, plant at least 1- $\frac{1}{4}$ inches but no more than 1- $\frac{1}{2}$ inches deep if moisture conditions are favorable. Check seed depth often in high-residue situations to make sure that seed is placed at the proper depth. These measurements should not include any surface residue. Seed left on the surface or in the residue layer will not grow or properly develop. If residue is problematic, consider residue management planter attachments.

Additional Information and References

Berg, R., D. DuBois, B. Jurgensen, R. Stevens, and G. Williamson. 2001. Date of planting corn. SE South Dakota Experiment Farm Progress Report 41:37-42. Accessed June 10, 2008. <http://plantsci.sdstate.edu>.

Carlson, C.G., T.A. Doerge, and D.E. Clay. 2002. Estimating corn yield losses from uneven spaced corn. SSMG-37 Clay et al. (ed.). Site-Specific Management Guidelines. Accessed June 10, 2008. <http://www.ppi-far.org/ssmg>.

Doerge, T.A. and T.E. Hall. 2000. The value of planter calibration using the MeterMax* System. Pioneer Hi-Bred International, Inc. Crop Insights 10:23. <http://www.pioneer.com>.

USDA-NASS. 2007. Annual Summary Bulletin. Accessed June 10, 2008. <http://www.nass.usda.gov/sd>.

Acknowledgements

Support for this document was provided by South Dakota State University, SDSU Extension, and the South Dakota Corn Utilization Council.



A G R O W I N G I N V E S T M E N T

Hall, R.G., K.D. Reitsma, and D.E. Clay. 2016. Appendix A: Corn Planting Guide. In Clay, D.E., C.G. Carlson, S.A. Clay, and E. Byamukama (eds). *iGrow Corn: Best Management Practices*. South Dakota State University.

The preceding is presented for informational purposes only. SDSU does not endorse the services, methods or products described herein, and makes no representations or warranties of any kind regarding them.

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at http://www.ascr.usda.gov/complaint_filing_cust.html and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by:

(1) mail: U.S. Department of Agriculture
Office of the Assistant Secretary for Civil Rights
1400 Independence Avenue, SW
Washington, D.C. 20250-9410;

(2) fax: (202) 690-7442; or

(3) email: program.intake@usda.gov.

USDA is an equal opportunity provider, employer, and lender.

SDSU Extension is an equal opportunity provider and employer in accordance with the nondiscrimination policies of South Dakota State University, the South Dakota Board of Regents and the United States Department of Agriculture.